

His-tag polyclonal antibody

Catalog: AP0032

Host: Rabbit

Reactivity: All

BackGround:

The H-H-H-H-H-H motif is used as a tag on many recombinant proteins to facilitate purification. A system that has proven to be very successful relies on the insertion of a six histidine (His6) sequence in the N-terminus of the encoded protein, allowing for efficient coupling to Ni²⁺-chelating resins and purification by single step affinity chromatography. This polyhistidine sequence can then be removed by specific cleavage at sites recognized by enzymes such as thrombin or enterokinase, permitting the separation of the target protein from the polyhistidine tag.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

N/A

Swiss-Prot:

N/A

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:2000~1:5000

IP: 1:50~1:200

IF: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:


His-tag polyclonal antibody detects over-expressed or recombinant proteins containing the 6×His epitope tag

DATA:




Western blot (WB) analysis of His-tag polyclonal antibody at 1:5000 dilution

Lane1:HEK293F whole cell lysate,transfected with pcDNA3.4-Flag-N-RBD-His.



Immunoprecipitation of 293F cell transfected with pcDNA3.4-Flag-N-RBD-His lysates using His-Tag pAb (Sepharose Bead Conjugate)#BD0048 (lane 3 and lane 4) and Nonspecific IgG Control (Sepharose Bead Conjugate)#BD0048 (lane 1 and lane 2) .Lane 5 is 30% input. The western blot was probed using His-Tag pAb #AP0032.



IF image of AP0032 stained HEK293F cells, transfected with pcDNA3.4-Flag-N-RBD-His. The cells were 4% paraformaldehyde fixed (20 min) and then incubated in 10% normal goat serum for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody His-tag pAb

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151



PRODUCT DATA SHEET

Bioworld Technology, Inc.

#AP0032(1:200) at 5 µg/ml overnight at +4 °C. The secondary antibody (Green) was Goat anti-Rabbit IgG (H+L) -FITC#BS10950. Hoechst33342 #BD5011 was used to stain the cell nuclei (blue).

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,
MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,
P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151