

ZAP70 (Tyr292) polyclonal antibody

Catalog: AP0158

Host: Rabbit

Reactivity: Human, Mouse

Background:

The activation of T lymphocytes by antigens is mediated by the T cell receptor (TCR), which is a multisubunit complex assembled from at least six different genes. The TCR subunits include the α and β chains, the CD3 γ , δ and ϵ chains and a ζ -containing homodimer or heterodimer. The protein tyrosine kinase ZAP-70 binds to the phosphorylated immunoreceptor tyrosine-base activation motifs (ITAMs) of the TCR ζ chain through two src-homology (SH2) domains. This binding results in the phosphorylation of ZAP-70 on multiple tyrosine residues, including Tyr292 and Tyr319. ZAP-70 is autophosphorylated on Tyr292, which is thought to negatively regulate ZAP-70 function in lymphocytes. Alternatively, ZAP-70 is positively regulated by phosphorylation on Tyr319, which mediates the SH2-dependent interaction between Lck and ZAP-70.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 70 kDa

Swiss-Prot:

P43403

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB:1:500~1:1000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

ZAP-70 (Tyr292) polyclonal antibody detects endogenous levels of ZAP-70 protein.

DATA:



Western blot (WB) analysis of ZAP70 (Tyr292) pAb

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151