PRODUCT DATA SHEET



Bioworld Technology,Inc.

Recombinant gAcrp30/Adipolean, Human

Catalog Number: BK0059-1mg Source: Escherichia coli. Quantity: 1mg

Description:

gAcrp30/Adipolean is the globular head domain of Adipocyte complement-related protein of 30 kDa (Acrp30), a cytokine expressed in adipocytes. The name of Acrp30 is bases on its closest homolog, complement factor c1q, and the globular domain of Acrp30 has an unexpected homolog with the Tumor Necrosis Factor (TNF) family of cytokines. Acrp30 is the isoform of adiponectin, and shares the two receptors with adiponectin: adipoR1 expressed in skeletal muscle, and adipoR2 expressed in liver. The expression level of Acrp30 in adipocytes is negatively correlated with body weight, and is lower in obese mouse than normal mouse. The globular domain of Acrp30 induces free fatty acid oxidation in muscle and weight reduction in mouse, therefore shows potential pharmacological effects in obesity. Recombinant human gAcrp30/Adipolean (rhgAcrp30) produced in E.coli is a single non-glycosylated polypeptide chain containing 144 amino acids. A fully biologically active molecule, rhgAcrp30 has a molecular mass of 16.6 kDa analyzed by reducing SDS-PAGE and is obtained by proprietary chromatographic techniques at GenScript.

Molecular Weight:

16.6 kDa, observed by reducing SDS-PAGE.

Purity

> 95% by SDS-PAGE analysis.

Biological Activity:

ED50 < 10 μ g/mL, measured by a cell growth inhibition assay using M1 cells, corresponding to a specific activity of > 100 units/mg.

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation:

Lyophilized after extensive dialysis against PBS.

AA Sequence:

KGEPGEGAYVYRSAFSVGLETYVTIPNM-PIRFTKIFYNQQNHYDGSTGKFHCNIPGLYY-FAYHITVYMKDVKVSLFKKDKAM-LFTYDQYQENNVDQASGSVLLHLEVGDQVWLQ VYGEGERNGLYADNDNDSTFTGFLLYHDTN

Endotoxin:

< 0.2 EU/µg, determined by LAL method.

Reconstitution:

Reconstituted in ddH2O at 100 µg/mL.

Storage:

Lyophilized recombinant human gAcrp30/Adipolean (rhgAcrp30) remains stable up to 6 months at -80 $^{\circ}$ C from date of receipt. Upon reconstitution, rhgAcrp30 should be stable up to 2 weeks at 4 $^{\circ}$ C or up to 3 months at -20 $^{\circ}$ C.

Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.