## PRODUCT DATA SHEET



Bioworld Technology,Inc.

# Recombinant MIP-3α/CCL20, Human

Catalog Number: BK0139-1mg Source: Escherichia coli. Quantity: 1mg

## **Description:**

Chemokine (C-C motif) ligand 20 (CCL20) or liver activation regulated chemokine (LARC) or Macrophage Inflammatory Protein-3 (MIP3A) is a small cytokine belonging to the CC chemokine family. It is strongly chemotactic for lymphocytes and weakly attracts neutrophils, also known as LARC (Liver and Activation-regulated Chemokine) and as Exodus. It can signal through the CCR6 receptor. MIP-3 alpha is chemotactic towards lymphocytes and dendritic cells. Additionally, it promotes the adhesion of memory CD4+ T cells and inhibits colony formation of bone marrow myeloid immature progenitors.Recombinant Human MIP-3α/CCL20 produced in E.coliis a single non-glycosylated polypeptide chain containing 70 amino acids. A fully biologically active molecule, rhMIP-3α/CCL20has a molecular mass of 8.0 kDa analvzed by reducing SDS-PAGE and is obtained by chromatographic techniques at GenScript.

### **Molecular Weight:**

8.0 kDa, observed by reducing SDS-PAGE.

#### **Purity:**

> 95% as analyzed by SDS-PAGE.

### **Biological Activity:**

The EC50 value of humanMIP-3 alpha/CCL20 on Ca^2+ mobilization assay in CHO-K1/G 15/hCCR6 cells (human Ga15 and human CCR6 stably expressed in CHO-K1 cells) is less than 200 ng/ml.

### **Physical Appearance:**

Sterile Filtered White lyophilized (freeze-dried) pow-

der.

#### **Formulation:**

Lyophilized after extensive dialysis against 53mM Na<sub>2</sub>HPO<sub>4</sub>, 147 mM NaH<sub>2</sub>PO<sub>4</sub>, 300 mM NaCl, pH6.5.

## **AA Sequence:**

ASNFDCCLGYT-DRILHPKFIVGFTRQLANEGCDINAIIFHT-KKKLSVCANPKQTWVKYIVRLLSKKVKNM

#### **Endotoxin:**

< 0.2 EU/µg, determined by LAL method.

### **Reconstitution:**

Reconstituted in ddH2O or PBS at 100 μg/ml.

## **Storage:**

Lyophilized recombinant Human MIP-  $3\alpha/CCL20$  remains stable up to 6 months at -80 °C from date of receipt. Upon reconstitution, recombinant human MIP- $3\alpha/CCL20$  should be stable up to 1 week at 4°C or up to 2 months at -20 °C.

### Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.

Email: <u>info@bioworlde.com</u> Tel: 6123263284 Fax: 6122933841 Tel: 0086-025-86371664 Fax:0086-025-86213570