

PRODUCT DATA SHEET



Bioworld Technology, Inc.

Recombinant IL-5, Mouse(CHO-expressed)

Catalog Number: BK0250-1mg

Source: CHO

Quantity: 1mg

Description:

Interleukin-5 (IL-5), produced by mast cells, T cells and eosinophils, is responsible for the activities attributed to eosinophil differentiating factor, B cell growth factor II and T cell-replacing factor (TRF). It can increase production and mobilization of eosinophils and CD34+ progenitors from the bone marrow. IL-5 plays an important role in inducing cell-mediated immunity against parasitic infections and certain tumors. IL-5 also promotes differentiation of basophils and primes them for histamine and leukotriene release. Recombinant Murine IL-5 is homodimeric protein with molecular weight ranging from 25 to 40 kDa due to glycosylation.

Molecular Weight:

25-40 kDa, observed by non-reducing SDS-PAGE.

Purity:

> 95% as analyzed by SDS-PAGE and HPLC.

Biological Activity:

ED50 < 0.5 ng/ml, measured in a cell proliferation assay using TF-1 cells, corresponding to a specific activity of > 2 × 10⁶ units/mg.

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation:

Lyophilized after extensive dialysis against PBS.

AA Sequence:

MEIPMSTVVKETLTQLSAHRALLTSNETMRL-
PVPTHKNHQLCIGEIFQGLDILK-
NQTVRGGTVEMLFQNLSLIKKYIDRQ
KEKCGEERRRTRQFLDYLQEFLGVMSTEW-
AMEG

Endotoxin:

< 0.2 EU/μg, determined by LAL method.

Reconstitution:

Reconstituted in ddH₂O or PBS at 100 μg/ml.

Storage:

Lyophilized recombinant Murine Interleukin 5 (IL-5) remains stable up to 6 months at -80 °C from date of receipt. Upon reconstitution, rmIL-5 should be stable up to 1 week at 4 °C or up to 2 months at -20 °C.

Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.