

# ATF2 (V65) polyclonal antibody

Catalog: BS1024

Host: Rabbit

Reactivity: Human, Mouse, Rat

## **BackGround:**

The transcription factor ATF-2 (also called CRE-BP1) binds to both AP-1 and CRE DNA response elements and is a member of the ATF/CREB family of leucine zipper proteins. ATF-2 interacts with a variety of viral oncoproteins and cellular tumor suppressors and is a target of the SAPK/JNK and p38 MAP kinase signaling pathways. Various forms of cellular stress, including genotoxic agents, inflammatory cytokines and UV irradiation, stimulate the transcriptional activity of ATF-2. Cellular stress activates ATF-2 by phosphorylation of Thr69 and Thr71. Both SAPK and p38 MAPK have been shown to phosphorylate ATF-2 at these sites in vitro and in cells transfected with ATF-2.

## **Product:**

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

### **Molecular Weight:**

~ 55 to 75 kDa

**Swiss-Prot:** 

#### P15336

#### **Purification&Purity:**

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

#### **Applications:**

WB: 1:500~1:1000

IHC: 1:50~1:200

IP: 1:50~1:200


## **Storage&Stability:**

Store at  $4 \,^{\circ}{\rm C}$  short term. Aliquot and store at  $-20 \,^{\circ}{\rm C}$  long term. Avoid freeze-thaw cycles.

## **Specificity:**

ATF2 (V65) polyclonal antibody detects endogenous levels of ATF2 protein.

#### **DATA:**




Western blot (WB) analysis of ATF2 (V65) polyclonal antibody at 1:500 dilution

Lane1:Hela cell lysate

Lane2:Mouse kidney tissue lysate

Lane3:Rat brain tissue lysate



Immunohistochemistry (IHC) analyzes of ATF2 (V65) pAb in paraf-

fin-embedded human breast carcinoma tissue at 1:100.

#### Note:

For research use only, not for use in diagnostic procedure.

## Bioworld Technology, Inc.

Add:	1660 South Highway 100, Suite 500 St. Louis Park,
	MN 55416,USA.
Email:	info@bioworlde.com
Tel:	6123263284
Fax:	6122933841

## Bioworld technology, co. Ltd.

 
 Add:
 No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

 Email:
 info@biogot.com

 Tel:
 0086-025-68037686

 Fax:
 0086-025-68035151