

Rad52 (N99) polyclonal antibody

Catalog: BS1691

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

DNA double-strand breaks are generated by ionizing radiation and endogenously produced radicals, and they often are repaired through the RAD52 homologous recombination pathway. The RAD52 family includes RAD51, RAD52, RAD54, RAD54B and MRE11 genes. Rad51 and Rad52 proteins perform the key steps in homologous recombination (HR), including the search for DNA homology and strand exchange, through similar mechanisms. Mre11 functions in both non-homologous end joining, and meiotic HR, and it is essential in mitosis for chromosome maintenance. Rad54 belongs to the SWI2/SNF2 subfamily of ATPases, which includes the DNA helicases involved in replication, recombination, and repair, as it contains seven amino acid sequence motifs that are largely conserved.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 46 kDa

Swiss-Prot:

P43351

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

IHC: 1:50~1:200

IF: 1:50~1:200

Storage&Stability:


Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Rad52 (N99) polyclonal antibody detects endogenous


levels of Rad52 protein.

DATA:



BS1691
Lot C136131

Immunohistochemistry (IHC) analyzes of Rad52 (N199) pAb in paraffin-embedded human lung carcinoma tissue at 1:50. showing Nucleus staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.



BS1691
Lot C136131

Immunohistochemistry (IHC) analyzes of Rad52 (N99) pAb in paraffin-embedded human stomach carcinoma tissue at 1:50. showing Nucleus staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151