

AIRE (P126) polyclonal antibody

Catalog: BS1879

Host: Rabbit

Reactivity: Human, Mouse, Rat

Background:

The autoimmune regulator gene, which is defective in the hereditary autoimmune disease APECED, encodes the transcriptional activator AIRE. AIRE is expressed in the medullary epithelial cells and monocyte-dendritic cells of the thymus, with lower expression in the spleen, fetal liver and lymph nodes. In adult tissue, AIRE expression in the thymus is confined to the medulla and the cortico-medullary junction, where it is modulated by thymocytes undergoing negative selection. At the cellular level, AIRE is located in microtubular structures of the cytoskeleton and in discrete nuclear dots resembling ND10 nuclear bodies. AIRE is induced by developing early thymocytes and is associated with the correct establishment of a regular thymic environment. AIRE regulates thymic architecture via transcriptional control of downstream target genes. AIRE mutations in APECED patients may affect thymic T cell selection and the formation of self-tolerance.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 60 kDa

Swiss-Prot:

O43918

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

IF: 1:50~1:200

Storage&Stability:


Store at 4 °C short term. Aliquot and store at -20 °C long

term. Avoid freeze-thaw cycles.

Specificity:

AIRE (P126) polyclonal antibody detects endogenous levels of AIRE protein.

DATA:




Western blot (WB) analysis of AIRE (P126) pAb at 1:500 dilution

Lane1:Panc1 whole cell lysate(40ug)

Lane2:A2780 whole cell lysate(40ug)


Lane3:The Heart tissue lysate of Rat(40ug)

Lane4:The Heart tissue lysate of Mouse(40ug)



BS1879
Lot CF36131

Immunohistochemistry (IHC) analyzes of AIRE (P126) pAb in paraffin-embedded human breast carcinoma tissue at 1:50, showing cytoplasmic and nucleus staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.



BS1879
Lot CF36131

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151



PRODUCT DATA SHEET

Bioworld Technology, Inc.

Immunohistochemistry (IHC) analyzes of AIRE (P126) pAb in paraffin-embedded human tonsil carcinoma tissue at 1:50, showing cytoplasmic and nucleus staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,
MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,
P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151