

ENX-2 (E192) polyclonal antibody

Catalog: BS2114

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

In Drosophila, the Polycomb (PcG) gene family encodes chromatin proteins that are required for the repression of homeotic loci in embryonic development. PcG proteins work in conjunction with the trithorax-group (trxG) proteins, which activate homeobox gene expression during embryonic development. ENX-1, a mammalian homolog of the Drosophila gene enhancer of zeste, is a PcG protein that is ubiquitously expressed during early embryogenesis and becomes restricted to the central and peripheral nervous systems and sites of fetal hematopoiesis during later development. In the adult, ENX-1 is restricted to specific sites, including spleen, testis and placenta. ENX-2 is another mammalian homolog of the Drosophila gene enhancer of zeste and contains one SET domain. The gene for human ENX-2 maps to chromosome 17q21. ENX-2 expression is ubiquitous in adult and fetal tissue, where it may aid in maintaining heterochromatin stability.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 85 kDa

Swiss-Prot:

Q92800

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

IF: 1:50~1:200

Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-20 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.


Bioworld Technology, Inc.

Add:	1660 South Highway 100, Suite 500 St. Louis Park,
	MN 55416,USA.
Email:	<u>info@bioworlde.com</u>
Tel:	6123263284
Fax:	6122933841

Specificity:


ENX-2 (E192) polyclonal antibody detects endogenous levels of ENX-2 protein.

DATA:



Western blot (WB) analysis of ENX-2 (E192) polyclonal antibody at 1:500 dilution

Lane1:MCF-7 whole cell lysate(40ug) Lane2:A2780 whole cell lysate(40ug) Lane3:HEK293T whole cell lysate(40ug) Lane4:A549 whole cell lysate(40ug) Lane5:HepG2 whole cell lysate(40ug) Lane6:H9C2 whole cell lysate(40ug) Lane7:3T3-L1 whole cell lysate(40ug)



Immunohistochemistry (IHC) analyzes of ENX-2 (E192) pAb in paraffin-embedded human brain tissue.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District N

 Add:
 No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

 Email:
 info@biogot.com

 Tel:
 0086-025-68037686

 Fax:
 0086-025-68035151



Bioworld Technology,Inc.

Bioworld Technology, Inc.		Biow	Bioworld technology, co. Ltd.	
Add:	1660 South Highway 100, Suite 500 St. Louis Park,	Add:	No 9, weidi road Qixia District Nanjing, 210046,	
	MN 55416,USA.		P. R. China.	
Email:	<u>info@bioworlde.com</u>	Email:	info@biogot.com	
Tel:	6123263284	Tel:	0086-025-68037686	
Fax:	6122933841	Fax:	0086-025-68035151	