

Catenin- δ 1 (G224) polyclonal antibody

Catalog: BS2992

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

The α , β , δ , and γ -catenins are proteins that bind to the highly conserved, intracellular cytoplasmic tail of E-cadherin. δ -catenin interacts with Presenilin 1 and is expressed in the brain. The gene encoding δ -catenin maps to human chromosome 5p15.2. A hemizygous loss of the gene encoding δ -catenin leads to the mental retardation associated with Cri-du-Chat syndrome. γ -catenin, also known as plakoglobin, is an 80-88 kDa protein that binds α -catenin and N-cadherin. In addition, the transmembrane phosphatase PTPm associates with catenin/cadherin complexes and may regulate complex signaling.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 108 kDa

Swiss-Prot:

O60716

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

IHC:1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Catenin- δ 1 (G224) polyclonal antibody detects endogenous levels of Catenin- δ 1 protein.

DATA:



Immunohistochemistry (IHC) analyzes of Catenin- δ 1 (G224) pAb in paraffin-embedded human tonsil carcinoma tissue at 1:50.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151