

CD63 (K156) polyclonal antibody

Catalog: BS3474

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

CD63 (also known as LAMP-3, Melanoma-associated antigen ME491, TSPAN30, MLA1 and OMA81H) is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 25-60 kDa

Swiss-Prot:

P08962

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:2000

IHC: 1:50~1:200


Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-20 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.


Specificity:

CD63 (K156) polyclonal antibody detects endogenous levels of CD63 protein.

DATA:



Western blot (WB) analysis of CD63 (K156) pAb at 1:1000 dilution Lane1:C6 whole cell lysate(30ug) Lane2:SP2/0 whole cell lysate(30ug) Lane3:K562 whole cell lysate(30ug) Lane4:A549 whole cell lysate(30ug)



Immunohistochemistry (IHC) analyzes of CD63 (K156) pAb

in paraffin-embedded human lung carcinoma tissue.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add:	1660 South Highway 100, Suite 500 St. Louis Park,
	MN 55416,USA.
Email:	info@bioworlde.com
Tel:	6123263284
Fax:	6122933841

Bioworld technology, co. Ltd.

 Add:
 No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

 Email:
 info@biogot.com

 Tel:
 0086-025-68037686

 Fax:
 0086-025-68035151