

# PRODUCT DATA SHEET

Bioworld Technology, Inc.

# ACAT1 (K266) polyclonal antibody

Catalog: BS3796 Host: Rabbit Reactivity: Human, Mouse, Rat

### **BackGround:**

ACAT-1 (acetyl-Coenzyme A acetyltransferase 1) is a mitochondrial enzyme involved in the formation and degradation of ketone bodies and is necessary for the proper metabolic processing of isoleucine. Rare defects in the gene encoding ACAT-1 lead to β-ketothiolase deficiency, which is characterized by ketoacidotic attacks. ACAT-2 (acetyl-Coenzyme A acetyltransferase 2) is considered a cytosolic protein and is crucial for cholesterol synthesis. Specifically, both Acetoacetyl-CoA specific thiolases, ACAT-1 and ACAT-2 catalyze the formation of acetoacetyl-CoA from two acetyl-CoA molecules. These enzymes are also capable of the reverse reaction, the cleavage of acetoacetyl-CoA into two acetyl-CoA molecules.

#### **Product:**

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

## **Molecular Weight:**

~ 45 kDa

# **Swiss-Prot:**

P24752

## **Purification&Purity:**

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

#### **Applications:**

WB: 1:500~1:1000 IHC: 1:50~1:200


## **Storage&Stability:**

Store at 4 ℃ short term. Aliquot and store at -20 ℃ long term. Avoid freeze-thaw cycles.

## **Specificity:**

ACAT1 (K266) polyclonal antibody detects endogenous levels of ACAT1 protein.

## **DATA:**




Western blot (WB) analysis of ACAT-1 (K266) pAb at 1:1000 dilution Lane1:HEK293T whole cell lysate(20ug)


Lane2:HepG2 whole cell lysate(20ug)

Lane3:The Lung tissue lysate of Rat(20ug)

Lane4: The Lung tissue lysate of Mouse(10ug)



Immunohistochemistry (IHC) analyzes of ACAT1 (K266) pA bin paraffin-embedded human liver carcinoma tissue at 1:50, showing cytoplasmic staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.



Immunohistochemistry (IHC) analyzes of ACAT1 (K266) pA bin paraffin-embedded human kidney carcinoma tissue at 1:50, showing cytoplasmic staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin fol-

## Bioworld Technology, Inc.

1660 South Highway 100, Suite 500 St. Louis Park, Add:

MN 55416,USA.

Email: info@bioworlde.com

Tel: 6123263284 6122933841 Fax:

# Bioworld technology, co. Ltd.

No 9, weidi road Qixia District Nanjing, 210046,

P. R. China.

Email: info@biogot.com Tel: 0086-025-68037686 0086-025-68035151 Fax:



# PRODUCT DATA SHEET

Bioworld Technology,Inc.

lowed by avidin-peroxidase.

For research use only, not for use in diagnostic procedure.

Note:

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,

MN 55416,USA.

Email: info@bioworlde.com

Tel: 6123263284 Fax: 6122933841 Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,

P. R. China.

Email: <u>info@biogot.com</u>
Tel: 0086-025-68037686
Fax: 0086-025-68035151