

AR (phospho-S650) polyclonal antibody

Catal	og:	BS4012
-------	-----	--------

Host:

Rabbit

Reactivity: Human,Rat

BackGround:

Androgens exhibit a wide range of effects on the development, maintenance and regulation of male phenotype and reproductive physiology in males. The androgen receptor (AR) is a member of the steroid superfamily of ligand-dependent transcription factors. ARs bind active testosterone (T) and dihydrotestosterone (DHT). The rates of association and dissociation of T are about 3 times more rapid than those of DHT. This difference in binding kinetics may account for the different physiological effects of T and DHT. Androgen binding results in an at least 6-fold increase in androgen receptor stability.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 110 kDa

Swiss-Prot:

P10275

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

IHC: 1:50~1:200


Storage&Stability:

Store at $4 \, \mathbb{C}$ short term. Aliquot and store at $-20 \, \mathbb{C}$ long term. Avoid freeze-thaw cycles.

Specificity:

p-AR (S650) polyclonal antibody detects endogenous levels of Androgen Receptor protein only when phosphorylated at Ser650.

DATA:



Immunohistochemistry (IHC) analyzes of p-AR (S650) pAb in paraffin-embedded human breast carcinoma tissue at 1:100.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA. **Email:** info@bioworlde.com Tel: 6123263284 6122933841 Fax:

Bioworld technology, co. Ltd. No 9, weidi road Qixia District Nanjing, 210046, Add: P. R. China. **Email:** info@biogot.com Tel: 0086-025-68037686 0086-025-68035151 Fax: