

HDAC10 Rabbit monoclonal antibody

Catalog: BS40311

Host: Rabbit

Reactivity: Human Mouse

BackGround:

Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2.

Molecular Weight:

75kDa

Swiss-Prot:

Q969S8

Purification&Purity:

Affinity-chromatography

Applications:

WB 1:1000~1:2000 ICC/IF 1:50~1:200 IP 1:50

Storage&Stability:


Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:


HDAC10 Antibody detects endogenous levels of

HDAC10

DATA:



Western blot analysis of HDAC10 expression in (1) HeLa cell lysate; (2) 3T3 cell lysate.



Immunofluorescent analysis of HeLa cells, using HDAC10 Antibody.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151