

NFκB-p65 (phospho-S468) polyclonal antibody

Catalog: BS4136

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

p65 is a subunit of the nuclear factor kappa B. The transcription factor NFκB is widely recognized as a critical mediator of immune and inflammatory responses. In most cell types, NFκB is found in the cytoplasm where it is associated with an inhibitory protein in many tissues. A high proportion of spontaneous NIH/3T3 transformants over-express c-Met and by transfection analysis the c-Met proto-oncogene has been shown to exhibit transforming activity.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 60, 75 kDa

Swiss-Prot:

Q04206

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

IHC: 1:50~1:200

IP: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

p-NFκB-p65 (S468) polyclonal antibody detects endogenous levels of NFκB-p65 protein when phosphorylated at Ser468.

DATA:



Immunohistochemistry (IHC) analyzes of p-NFκB-65 (S276) pAb in paraffin-embedded human breast carcinoma tissue at 1:100.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151