

PRODUCT DATA SHEET

Bioworld Technology,Inc.

HDAC4 (phospho-S632) polyclonal antibody

Catalog: BS4233 Host: Rabbit Reactivity: Human, Mouse, Rat

BackGround:

In the intact cell, DNA closely associates with histones and other nuclear proteins to form chromatin. The remodeling of chromatin is believed to be a critical component of transcriptional regulation, and a major source of this remodeling is brought about by the acetylation of nucleosomal histones. Acetylation of lysine residues in the amino-terminal tail domain of histone results in an allosteric change in the nucleosomal conformation and an increased accessibility to transcription factors by DNA. Conversely, the deacetylation of histones is associated with transcriptional silencing. Several mammalian proteins have been identified as nuclear histone acetylases, including GCN5, p300/CBP, PCAF (p300/CBPassociated factor), HAT1 and the TFIID subunit TAF II p250. Mammalian HDAC1 (also designated HD1), HDAC2 (also designated RPD3) and HDAC3-6 have been identified as histone deacetylases.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 119 kDa

Swiss-Prot:

P56524

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific im-

munogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at $4\,\mathrm{C}$ short term. Aliquot and store at -20 C long term. Avoid freeze-thaw cycles.

Specificity:

p-HDAC4 (S632) polyclonal antibody detects endogenous levels of HDAC4 protein only when phosphorylated at Ser632.

DATA:



Western blot (WB) analysis of p-HDAC4 (S632) polyclonal antibody in extracts from Jurkat cells.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,

MN 55416,USA.

Email: <u>info@bioworlde.com</u>

Tel: 6123263284 Fax: 6122933841 Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,

P. R. China.

Email: <u>info@biogot.com</u>
Tel: 0086-025-68037686
Fax: 0086-025-68035151