

eIF2 α (phospho-S51) polyclonal antibody

Catalog: BS4787

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

eIF2 α , also known as EIF2S1 or EIF2, is a 315 amino acid subunit of the eukaryotic initiation complex that functions to bind tRNA to the 40S ribosomal subunit (in a GTP-dependent manner), thereby initiating translation. In addition, the phosphorylation state of eIF2 α controls the rate of tRNA translation. When eIF2 α is not phosphorylated, translation occurs at a normal rate. However, upon phosphorylation by one of several kinases, eIF2 α is stabilized, thus preventing the GDP/GTP exchange reaction and slowing translation.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 38 kDa

Swiss-Prot:

P05198

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

Storage&Stability:


Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

p-eIF2 α (S51) polyclonal antibody detects endogenous levels of eIF2 α protein only when phosphorylated at

Ser51.

DATA:



Western blot (WB) analysis of p-eIF2 α (S51) pAb at 1:500 dilution

Lane1:HeLa whole cell lysate(40ug)


Lane2:MCF-7 whole cell lysate(40ug)

Lane3:The Uterus tissue lysate of Rat(40ug)

Lane4:The Lung tissue lysate of Mouse(40ug)

Lane5:A2780 whole cell lysate(40ug)

Lane6:H1792 whole cell lysate(40ug)



Immunohistochemistry (IHC) analyzes of p-eIF2 α (S51) pAb in paraffin-embedded human breast carcinoma tissue.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151