

PRODUCT DATA SHEET



Bioworld Technology, Inc.

4E-BP1 (P71) pAb

Cat No.: BS5501

Host: Rabbit

Reactivity: Human, Mouse, Rat

BACKGROUND

The translation of proteins from eukaryotic mRNA is initiated by the multisubunit complex eIF-4F, which associates with the mRNA 5' cap structure. eIF-4E, a component of eIF-4F, is responsible for binding to the 5' cap structure and for the assembly of the eIF-4F complex. The regulatory protein 4E-BP1, also referred to as PHAS-I, inhibits eIF-4E function. Phosphorylation of 4E-BP1 by S6 kinase p70, MAP kinases or PKCs causes the disassociation of 4E-BP1 from eIF-4E, promoting translation. A protein that is functionally related to 4E-BP1, designated 4E-BP2, also associates with eIF-4E.

PRODUCT

1 mg/ml in Phosphate buffered saline (PBS) with 0.05

PURIFICATION & PURITY

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

APPLICATIONS

IHC: 1:50 ~ 1:200 (Recommended Dilutions)


STORAGE & STABILITY

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.

SPECIFICITY

4E-BP1 (P71) pAb detects endogenous levels of 4E-BP1 protein.

DATA



Immunohistochemistry (IHC) analyzes of 4E-BP1 (P71) pAb in paraffin-embedded human breast carcinoma tissue.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

Bioworld Technology, Inc.

1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA. Tel: 6123263284
www.bioworld.com Orders: order@bioworld.com Support: support@bioworld.com