

SLP-76 polyclonal antibody

Catalog: BS5935

Host: Ra

Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Overexpression of Vav or SLP-76 in Jurkat cells leads to NFAT activation and IL-2 production. When co-expressed, Vav and SLP-76 synergize to induce a robust basal and TCR-mediated IL-2 response. Although SLP-76 does not contain a motif that would indicate it to be a member of the tyrosine, serine/threonine or lipid kinase families, it does contain several putative SH2/SH3-binding domains and has been shown to physically associate with the adapter protein GRB2 as well as PLC g1. The discovery of SLP-76 represents an important step in elucidating the mechanism of Vav transformation and TCR-mediated NFAT activation.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 75 kDa

Swiss-Prot:

Q13094

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-20 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.

Specificity:

SLP-76 polyclonal antibody detects endogenous levels of SLP-76 protein.

DATA:



Western blot (WB) analysis of SLP-76 pAb at 1:1000 dilution Lane1:Raw264.7 whole cell lysate(40ug) Lane2:The Thymus tissue lysate of Mouse(40ug)

Lane3: The Thymus tissue lysate of Rat(40ug)

Lane4:K562 whole cell lysate(40ug)

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

 Add:
 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.

 Email:
 info@bioworlde.com

 Tel:
 6123263284

 Fax:
 6122933841

Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151