

Endoglin polyclonal antibody

Catalog: BS60010

Host: 1

Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Hereditary hemorrhagic telangiectasia (HHT) is an autosomal dominant disorder characterized by vascular abnormalities such as dilated vessels, hemorrhages, liver and lung congestion, and brain or heart ischemia. Mutations in two genes, Endoglin (also designated CD105) and ALK-1 (Activin receptorlike kinase 1, also designated TGF^β superfamily RI), are responsible for HHT. Endoglin is mutated in HHT1, and ALK-1 is mutated in HHT2, both of which are thought to be caused by haploinsufficiency. Endoglin and ALK-1 are type III and type I members of the TGFβ receptor superfamily, respectively, that are expressed on vascular endothelial cells. Endoglin can only bind ligands of the TGFB superfamily via association with the respective ligand binding receptors for TGF_{β1}, TGF_{β3}, Activin-A, BMP-2 and BMP-7. The human ALK-1 gene encodes two protein species which exist as a result of either glycosylation or alternative splicing events. ALK-1 preferentially binds TGFβ1 and is expressed in bone marrow stromal cells, lung, brain, kidney and spleen.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 100 kDa

Swiss-Prot:

P17813

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific im-

munogen and the purity is > 95% (by SDS-PAGE).

Applications: WB: 1:500~1:1000

Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-20 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.

Specificity:

Endoglin polyclonal antibody detects endogenous levels of Endoglin protein.

DATA:



Western blot (WB) analysis of ENG pAb at 1:500 dilution Lane1:PC3 whole cell lysate(40ug) Lane2:HEK293T whole cell lysate(40ug) Lane3:Hela whole cell lysate(40ug) Lane4:PC12 whole cell lysate(40ug) Lane5:3T3-L1 whole cell lysate(40ug)

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

 Add:
 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.

 Email:
 info@bioworlde.com

 Tel:
 6123263284

 Fax:
 6122933841

Bioworld technology, co. Ltd.

 Add:
 No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

 Email:
 info@biogot.com

 Tel:
 0086-025-68037686

 Fax:
 0086-025-68035151