

HNF-4 α / γ polyclonal antibody

Catalog: BS61772

Host: Rabbit

Reactivity: Human, Mouse, Rat

Background:

Hepatocyte nuclear factor 4 α (HNF4 α) is a transcription factor that belongs to the steroid hormone receptor superfamily and is enriched in liver. HNF4 α , in association with PGC-1 α , activates gluconeogenic genes such as phosphoenolpyruvate carboxykinase and glucose-6-phosphatase genes in fasted livers. Conditional knockout of the HNF4 α gene in the mouse liver destroys lipid homeostasis and leads to lipid accumulation in the liver and a reduction of serum cholesterol and triglyceride levels. Mutations in HNF4 α have been linked to maturity-onset diabetes of the young (MODY).

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 52 kDa

Swiss-Prot:

P41235/Q14541

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

HNF-4 α / γ polyclonal antibody detects endogenous levels of HNF-4 α / γ protein.

DATA:



Western blot (WB) analysis of HNF-4 α / γ polyclonal antibody at 1:500 dilution

Lane1:HCT116 whole cell lysate(40ug)

Lane2:K562 whole cell lysate(40ug)

Lane3:Hela whole cell lysate(40ug)

Lane4:CT26 whole cell lysate(40ug)

Lane5:A549 whole cell lysate(40ug)

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151