

Androgen Receptor polyclonal antibody

Catalog: BS62258

Host: Rabbit

Reactivity: Human, Rat

BackGround:

Androgens exhibit a wide range of effects on the development, maintenance and regulation of male phenotype and reproductive physiology in males. The androgen receptor (AR) is a member of the steroid superfamily of ligand-dependent transcription factors. ARs bind active testosterone (T) and dihydrotestosterone (DHT). The rates of association and dissociation of T are about 3 times more rapid than those of DHT. This difference in binding kinetics may account for the different physiological effects of T and DHT. Androgen binding results in an at least 6-fold increase in androgen receptor stability.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 110 kDa

Swiss-Prot:

P10275

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB:1:500~1:1000


Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-20 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.

Specificity:

Androgen Receptor polyclonal antibody detects endogenous levels of Androgen Receptor protein.

DATA:



Western blot (WB) analysis of Androgen Receptor polyclonal antibody at 1:500 dilution

LaneA:Hela whole cell lysate

LaneB:MCF-7 whole cell lysate

LaneC: The Testis tissue lysate of Rat

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

 Add:
 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.

 Email:
 info@bioworlde.com

 Tel:
 6123263284

 Fax:
 6122933841

Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: info@biogot.com Tel: 0086-025-68037686 Fax: 0086-025-68035151