

SIRT1 polyclonal antibody

Catalog: BS64501

Host: Rabbit

Reactivity: Human

BackGround:

The Silent Information Regulator (SIR2) family of genes is a highly conserved group of genes that encode nicotinamide adenine dinucleotide (NAD)-dependent protein deacetylases, also known as class III histone deacetylases. The first discovered and best characterized of these genes is *Saccharomyces cerevisiae* SIR2, which is involved in silencing of mating type loci, telomere maintenance, DNA damage response, and cell aging. SirT1, the mammalian ortholog of Sir2, is a nuclear protein implicated in the regulation of many cellular processes, including apoptosis, cellular senescence, endocrine signaling, glucose homeostasis, aging, and longevity. Targets of SirT1 include acetylated p53, p300, Ku70, forkhead (FoxO) transcription factors, PPAR γ , and the PPAR γ coactivator-1 α (PGC-1 α) protein (8). Deacetylation of p53 and FoxO transcription factors represses apoptosis and increases cell survival. Deacetylation of PPAR γ and PGC-1 α regulates the gluconeogenic/glycolytic pathways in the liver and fat mobilization in white adipocytes in response to fasting. SirT1 deacetylase activity is inhibited by nicotinamide and activated by resveratrol. In addition, SirT1 activity may be regulated by phosphorylation, as it is phosphorylated at Ser27 and Ser47 *in vivo*; however, the function of these phosphorylation sites has not yet been determined.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2.

Molecular Weight:

~120kDa

Swiss-Prot:

Q96EB6

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:2000~1:5000

Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

SIRT1 polyclonal antibody detects endogenous levels of SIRT1 protein.

DATA:



Western blot (WB) analysis of SIRT1 polyclonal antibody at 1:2000 dilution

Lane1:HEK293T whole cell lysate(40ug)

Lane2:HCT116 whole cell lysate(20ug)

Lane3:Hela whole cell lysate(40ug)

Lane4:CT-26 whole cell lysate(40ug)

Lane5:C6 whole cell lysate(40ug)

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151