

Rel B (S552) polyclonal antibody

Catalog: BS70007

Host: Rabbit

Reactivity: Human, Mouse

Background:

The NFκB transcription factor was originally identified as a protein complex consisting of a DNA-binding subunit and an associated protein. The DNA-binding subunit is functionally related to c-Rel p75 and Rel B p68. The p50 subunit was initially believed to be a functionally unique protein derived from the amino-terminus of a precursor designated p105. A second protein designated p52 (previously referred to as p49) has been identified that can act as an alternative NFκB subunit. Rel B does not bind with high affinity to NFκB sites, but heterodimers between Rel B and p50 bind with an affinity comparable to that of p50 NFκB homodimers. However, Rel B/p50 heterodimers, in contrast to NFκB heterodimers, transactivates transcription of promoters containing κB binding sites.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 70 kDa

Swiss-Prot:

Q01201

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific im-

munogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:2000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Rel B (S552) polyclonal antibody detects endogenous levels of Rel B protein.

DATA:



Western blot analysis of Rel B expression in HepG2 cell lysates. The lane on the left is treated with the antigen-specific peptide.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151