

IKK α / β polyclonal antibody

Catalog: BS90697

Host: Rabbit

Reactivity: Human, Mouse, Rat

Background:

The transcription factor NF κ B is retained in the cytoplasm in an inactive form by the inhibitory protein I κ B. Activation of NF κ B requires that I κ B be phosphorylated on specific serine residues, which results in targeted degradation of I κ B. I κ B kinase a (IKK α), previously designated CHUK, interacts with I κ B- α and specifically phosphorylates I κ B- α on Ser 32 and 36, the sites that trigger its degradation. IKK α appears to be critical for NF κ B activation in response to proinflammatory cytokines. Phosphorylation of I κ B by IKK α is stimulated by the NF κ B inducing kinase (NIK), which itself is a central regulator for NF κ B activation in response to TNF and IL-1. The functional IKK complex contains three subunits, IKK α , IKK β and IKK γ (also designated NEMO), and each appear to make essential contributions to I κ B phosphorylation.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

85/87 kDa

Swiss-Prot:

O14920(Human) O15111(Human) O88351(Mouse)
Q60680(Mouse) Q9QY78(Rat)

Purification&Purity:

ProA affinity purified

Applications:

WB:1:1,000

ICC:1:100-1:500

Storage&Stability:


Store at +4 °C after thawing. Aliquot store at -20 °C or

-80 °C. Avoid repeated freeze / thaw cycles.

Specificity:


IKK α / β polyclonal antibody detects endogenous levels of IKK α / β protein.

DATA:



Western blot analysis of IKK alpha+IKK beta on different lysates using anti-IKK alpha+IKK beta antibody at 1/1,000 dilution. Positive control:

Lane 1: Hela Lane 2: Daudi Lane 3: A431



ICC staining IKK alpha+IKK beta in HeLa cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151