Raptor polyclonal antibody

Catalog: BS91154 Host: Rabbit Reactivity: Human, Mouse, Rat

BackGround:

Regulatory associated protein of FRAP, also designated Raptor, is a binding partner for mammalian target of rapamycin kinase (FRAP), and is essential for FRAP signalling in vivo. Raptor binding to FRAP is critical for FRAP-catalysed substrate phosphorylation of 4E-BP1. The raptor-FRAP complex is nutrient-sensitive and is important for a mechanism by which cells coordinate cell growth and size with changing environmental conditions. Raptor serves as a negative regulator of FRAP kinase activity under nutrient-deprived conditions and is an important component in the FRAP pathway. Raptor is highly expressed in skeletal muscle and to a lesser extent in brain, kidney, lung and placenta.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

131 kDa

Swiss-Prot:

Q8N122(Human) Q8K4Q0(Mouse)

Purification&Purity:

Peptide affinity purified.

Applications:

WB:1:500

ICC:1:50-1:200

IHC:1:50-1:200

FC:1:50-1:100

Storage&Stability:

Store at +4℃ after thawing. Aliquot store at -20℃. Avoid repeated freeze / thaw cycles.

Specificity:

Raptor polyclonal antibody detects endogenous levels of Raptor protein.

DATA:



Western blot analysis of Raptor on MCF-7 cell lysate using anti-Raptor antibody at 1/500 dilution.



ICC staining Raptor in 293T cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.
Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.
Email: info@bioworlde.com
Tel: 6123263284
Fax: 6122933841

Bioworld technology, co. Ltd.
Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.
Email: info@biogot.com
Tel: 0086-025-68037686
Fax: 0086-025-68035151