

AMPK β 1 (V175) polyclonal antibody

Catalog: BS9234

Host: Rabbit

Reactivity: Human, Mouse, Rat

Background:

AMPK β 1 is a member of the SNF1 kinase family. It is a heterotrimeric protein comprising of an alpha (catalytic), beta (non-catalytic) and gamma (non-catalytic) subunits, 63, 38 and 38kDa respectively. AMPK β 1 regulates fatty acid and sterol synthesis by phosphorylation of acetyl-CoA. It also regulates cholesterol synthesis via phosphorylation and in-activation of hydroxymethylglutaryl-CoA reductase. Two isoforms of AMPK β 1 have been found.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 30, 38 kDa

Swiss-Prot:

Q9Y478

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

IHC: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:


AMPK β 1 (V175) polyclonal antibody detects endogenous levels of AMPK β 1 protein.

DATA:




BS9234
Lot CA36131

Immunohistochemistry (IHC) analyzes of AMPK β 1 (V175) pAb in paraffin-embedded human breast carcinoma tissue at 1:50, showing cytoplasmic and nuclear staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.



BS9234
Lot CA36131

Immunohistochemistry (IHC) analyzes of AMPK β 1 (V175) pAb in paraffin-embedded human colon carcinoma tissue at 1:50, showing cytoplasmic and nuclear staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.



BS9234
Lot CA36131

Immunohistochemistry (IHC) analyzes of AMPK β 1 (V175) pAb in paraffin-embedded human lung carcinoma tissue at 1:50, showing cytoplasmic and nuclear staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151



PRODUCT DATA SHEET

Bioworld Technology, Inc.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,
MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,
P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151