

ADAM32 (N530) polyclonal antibody

| Catalog: | BS9290 | Host: | Rabbit | Reactivity: | Human,Mouse,Rat |
|---|----------|-------|--------------|--|-----------------|
| BackGround: | | | | munogen and the purity is $> 95\%$ (by SDS-PAGE). | |
| ADAM (disintegrin and metalloproteinase domain) pro- | | | | Applications: | |
| teins, also known as MDC (metalloproteinase, disintegrin | | | | WB: 1:500~1:1000 | |
| | | | | | |
| and cysteine-rich domain) proteins or cellular disinteg- | | | | Storage&Stability: | |
| rins, are a family of proteins that are expressed in nu- | | | | Store at 4 ${}^{\ensuremath{\mathbb{C}}}$ short term. Aliquot and store at -20 ${}^{\ensuremath{\mathbb{C}}}$ long | |
| merous tissues. ADAMs are membrane-anchored, glyco- | | | | term. Avoid freeze-thaw cycles. | |
| sylated, Zn2+ dependent proteases that catalyze proteoly- | | | | Specificity: | |
| sis, adhesion, fusion and intracellular signaling. The | | | | ADAM32 (N530) polyclonal antibody detects endoge- | |
| ADAM family consists of more than 30 different mem- | | | | nous levels of ADAM32 protein. | |
| bers with many diverse functions. ADAM32 is expressed | | | | DATA: | |
| predominantly in the testis on the sperm surface. The | | | | | |
| ADAM32 precursor originates in the testis and is pro- | | | and is pro- | Western blot (WB) analysis of ADAM32 (N530) polyclonal antibody at 1:500 dilution Lane1:BV2 whole cell lysate(40ug) Lane2:The Testis tissue lysate of Rat(40ug) | |
| cessed during epididymal maturation. ADAM32 may play | | | 32 may play | | |
| a role in sperm-egg adhesion or sperm development. | | | pment. | | |
| Product: | | | | | |
| Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, | | | odium azide, | Lane3:MCF-7 whole cell lysate(40ug) | |
| 50% glycerol, pH7.2 | | | | Lane4:Hela whole cell lysate(20ug) | |
| Molecular Weight: | | | | Lane5:U-87MG whole cell lysate(30ug) Note: | |
| ~ 87 kDa | ~ 87 kDa | | | | |
| Swiss-Prot: | | | | For research use only, not for use in diagnostic procedure. | |
| Q8TC27 | | | | | |
| Purification&Purity: | | | | | |

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific im-

Bioworld Technology, Inc. Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA. Email: <u>info@bioworlde.com</u> Tel: 6123263284 Fax: 6122933841

Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: <u>info@biogot.com</u> Tel: 0086-025-68037686 Fax: 0086-025-68035151