

Smad2 (Phospho-S255) polyclonal antibody

Catalog: BS94015

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Smad proteins, the mammalian homologs of the Drosophila mothers against decapentaplegic (Mad), have been implicated as downstream effectors of TGF β /BMP signaling. Smad1 (also designated Madr1 or JV4-1) and Smad5 are effectors of BMP-2 and BMP-4 function, while Smad2 (also designated Madr2 or JV18-1) and Smad3 are involved in TGF β and Activin-mediated growth modulation. Smad4 (also designated DPC4) has been shown to mediate all of the above activities through interaction with various Smad family members. Smad6 and Smad7 regulate the response to Activin/TGF β signaling by interfering with TGF β -mediated phosphorylation of other Smad proteins.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

52 kDa

Swiss-Prot:

Q15796(Human) Q62432(Mouse) O70436(Rat)

Purification&Purity:

ProA affinity purified

Applications:

WB:1:1,000

IHC:1:50-1:200


Storage&Stability:

Store at +4 °C after thawing. Aliquot store at -20 °C or -80 °C. Avoid repeated freeze / thaw cycles.

Specificity:

Smad2 (Phospho-S255) polyclonal antibody detects endogenous levels of Smad2 protein only when phosphorylated at S255.

DATA:



Immunohistochemical analysis of paraffin-embedded mouse uterus tissue using anti-Phospho-Smad2(S255) antibody. Counter stained with hematoxylin.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151