

MKK2 monoclonal antibody


Catalog:	MB65962	Host:	Mouse	Reactivity:	Human, Mouse, Rat
BackGround:				Applications:	
Catalyzes the concomitant phosphorylation of a threonine				WB (1/1000 - 1/2000)	
and a tyrosine residue in a Thr-Glu-Tyr sequence located				Storage&Stability:	

Store at $4 \,^{\circ}$ short term. Aliquot and store at $-20 \,^{\circ}$ long term. Avoid freeze-thaw cycles.

Specificity:

Recognizes endogenous levels of MKK2 protein.

DATA:



Western blot analysis of MKK2 expression in Hela (A), NIH3T3 (B), rat brain (C) whole cell lysates.

Note:

For research use only, not for use in diagnostic procedure.

Catalyzes the concomitant phosphorylation of a threonine and a tyrosine residue in a Thr-Glu-Tyr sequence located in MAP kinases. Activates the ERK1 and ERK2 MAP kinases (By similarity).

Activates BRAF in a KSR1 or KSR2-dependent manner; by binding to KSR1 or KSR2 releases the inhibitory intramolecular interaction between KSR1 or KSR2 protein kinase and N-terminal domains which promotes KSR1 or KSR2-BRAF dimerization and BRAF activation (Pub-Med:29433126).

Product:

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

Molecular Weight:

~ 44 kDa

Swiss-Prot:

P36507

Purification&Purity:

The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Bioworld Technology, Inc. Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA. Email: info@bioworlde.com Tel: 6123263284 Fax: 6122933841

Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: info@biogot.com Tel: 0086-025-68037686 Fax: 0086-025-68035151