

# **AKT Rabbit monoclonal antibody**

Catalog: MB66248 Host:

Rabbit

Reactivity: Human, Mouse, Rat, Hamster

## **BackGround:**

Akt, also referred to as PKB or Rac, plays a critical role in controlling survival and apoptosis. This protein kinase is activated by insulin and various growth and survival factors to function in a wortmannin-sensitive pathway involving PI3 kinase. Akt is activated by phospholipid binding and activation loop phosphorylation at Thr308 by PDK1 and by phosphorylation within the carboxy terminus at Ser473. The previously elusive PDK2 responsible for phosphorylation of Akt at Ser473 has been identified as mammalian target of rapamycin (mTOR) in a rapamycin-insensitive complex with rictor and Sin1. Akt promotes cell survival by inhibiting apoptosis through phosphorylation and inactivation of several targets, including Bad, forkhead transcription factors, c-Raf, and caspase-9. PTEN phosphatase is a major negative regulator of the PI3K/Akt signaling pathway. LY294002 is a specific PI3 kinase inhibitor. Another essential Akt function is the regulation of glycogen synthesis through phosphorylation and inactivation of GSK-3 $\alpha$  and  $\beta$ . Akt may also play a role in insulin stimulation of glucose transport. In addition to its role in survival and glycogen synthesis, Akt is involved in cell cycle regulation by preventing GSK-3\beta-mediated phosphorylation and degradation of cyclin D1 and by negatively regulating the cyclin-dependent kinase inhibitors p27 Kip1 and p21 Waf1/Cip1. Akt also plays a critical role in cell growth by directly phosphorylating mTOR in a rapamycin-sensitive complex containing raptor. More importantly, Akt phosphorylates and inactivates tuberin (TSC2), an inhibitor of mTOR within the mTOR-raptor complex.

#### **Product:**

Liquid in 50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 50% Glycerol, 0.01% Sodium azide and 0.05% BSA.

**Molecular Weight:** 

~ 55 kDa

### **Bioworld Technology, Inc.**

Add:	1660 South Highway 100, Suite 500 St. Louis Park,
	MN 55416,USA.
Email:	info@bioworlde.com
Tel:	6123263284
Fax:	6122933841

# **Swiss-Prot:**

# P31749; P31751; Q9Y243

# **Purification&Purity:**

The antibody was purified by immunogen affinity chromatography.

## **Applications:**

WB (1/500 - 1/1000), IHC (1/50 - 1/100), IF/ICC (1/50 -1/100), IP (1/10 - 1/50)

#### **Storage&Stability:**

k

Store at  $4 \, \mathbb{C}$  short term. Aliquot and store at  $-20 \, \mathbb{C}$  long term. Avoid freeze-thaw cycles.


#### **Specificity:**

Recognizes endogenous levels of AKT protein.

#### **DATA:**

Da	А	в	С	D
130 95				
72				
55	-	-	-	-
43				
34				
26				
17				

Western blot analysis of AKT expression in Hela (A), Jurkat (B), rat brain (C), CHO-K1 (D) whole cell lysates.



Immunohistochemical analysis of AKT staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then


Bioworld technology, co. Ltd. No 9, weidi road Qixia District Nanjing, 210046, Add: P. R. China. **Email:** info@biogot.com Tel: 0086-025-68037686 0086-025-68035151 Fax:



# **PRODUCT DATA SHEET**

## **Bioworld Technology,Inc.**

counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of AKT staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark.

#### Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.			Bioworld technology, co. Ltd.		
Add:	1660 South Highway 100, Suite 500 St. Louis Park,	Add:	No 9, weidi road Qixia District Nanjing, 210046,		
	MN 55416,USA.		P. R. China.		
Email:	<u>info@bioworlde.com</u>	Email:	info@biogot.com		
Tel:	6123263284	Tel:	0086-025-68037686		
Fax:	6122933841	Fax:	0086-025-68035151		