

Catenin- β Mouse monoclonal antibody

Catalog: MB9005

Host: Mouse

Reactivity: Human,mouse,rat

BackGround:

β -Catenin is a key downstream effector in the Wnt signaling pathway. It is implicated in two major biological processes in vertebrates: early embryonic development and tumorigenesis. CK1 phosphorylates β -catenin at Ser45. This phosphorylation event primes β -catenin for subsequent phosphorylation by GSK-3 β . GSK-3 β destabilizes β -catenin by phosphorylating it at Ser33, Ser37, and Thr41. Mutations at these sites result in the stabilization of β -catenin protein levels and have been found in many tumor cell lines.

Product:

Mouse IgG1, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 92 kDa

Swiss-Prot:

P35222

Purification&Purity:

Protein A affinity purified

Applications:

WB: 1:2000-1:5000

ICC: 1:100-1:200

FC: 1:50


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

This antibody detects endogenous levels of Catenin- β and does not cross-react with related proteins.

DATA:



Western blot (WB) analysis of Catenin- β Mouse mAb at 1:2000 dilution


Lane1:A431 whole cell lysate

Lane2:NCCIT whole cell lysate

Lane3:SW480 whole cell lysate

Lane4:HT-29 whole cell lysate

Lane5:The brain tissue lysate of Mouse



ICC staining Catenin- β in NCCIT cells (red). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton *100/PBS.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151