

CD82 Recombinant Protein

Catalog:	NCP0231	Host:	E.coli	Tag:	His-tag
----------	---------	-------	--------	------	---------

BackGround:

CD82 (KAI1) belongs to the tetraspanin family, which is characterized by four transmembrane domains, one short extracellular domain (ECL1), and one long extracellular domain (ECL2). CD82 does not have enzymatic activity and appears to function by regulating the trafficking of other proteins and organization of the cell membrane. CD82 was originally described as a costimulator for T cells that directly associates with CD4 and CD8, and was subsequently identified during a screen as a metastasis suppressor in prostate cancer. CD82 has since been found to act as a metastasis suppressor in a variety of cancers, and its downregulation is associated with poor prognosis in research studies. CD82 suppresses metastasis through multiple mechanisms including inhibition of cell motility and invasion by modulating c-Met and the urokinase plasminogen activator surface receptor (uPAR), as well as promotion of homotypic cell-cell adhesion by stabilizing interactions between E-cadherin and β-catenin.

Product:

PBS, 4M Urea, PH7.4

Molecular Weight:

~15kDa

Swiss-Prot:

P27701

Purification&Purity:

Transferred into competent cells and the supernatant was purified by NI column affinity chromatography and the purity is > 85% (by SDS-PAGE).

Restriction Sites:

NdeI-XhoI

Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-20 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.

Expression Vector:

pet-22b(+)

DATA:

	M	S
KDa 116		
66.2		
45	1000	月 1 2)
35		
25		
18.4	_	
14.4		-

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.				
Add:	1660 South Highway 100, Suite 500 St. Louis Park,			
	MN 55416,USA.			
Email:	info@bioworlde.com			
Tel:	6123263284			
Fax:	6122933841			