

Bioworld Technology, Inc.

Recombinant Human Ciliary Neurotrophic Factor (rHuCNTF)

Catalog Number: PR1012 Source: Escherichia coli. Quantity:5µg/20µg/1.0mg

Description

Ciliary neurotrophic factor (CNTF) is a polypeptide initially purified from chick embryo ocular tissue and identified as a trophic factor for embryonic chick ciliary parasympathetic neurons in culture. Subsequent studies have demonstrated that CNTF is a survival factor for additional neuronal cell types including: dorsal root ganglion sensory neurons, sympathetic ganglion neurons, embryonic motor neurons, major pelvic ganglion neurons and hippocampal neurons. CNTF has also been shown to prevent the degeneration of motor axons after axotomy.

The gene for human CNTF has been localized to the proximal region of the long arm of chromosome 11. The cDNA for human CNTF encodes a 200 amino acid residue polypeptide that lacks a signal sequence. CNTF is highly conserved across species and exhibits cross-species activities. Human and rat CNTF share approximately 83% homology in their protein sequence. CNTF is structurally related to IL-6, IL-11, LIF and OSM. All of these four helix bundle cytokines share gp130 as a signal-transducing subunit in their receptor complexes.

Molecular Weight:

Approximately 22.8 kDa, a single non-glycosylated polypeptide chain containing 199 amino acids.

Purity:

>97% by SDS-PAGE and HPLC analyses.

Biological Activity:

ED50 as determined by the dose-dependant stimulation of kept at -20 \(\text{C} \) for long term storage, preferably desiccated.

TF-1 cells is less than 2.0 ng/ml, corresponding to a specific activity of $> 5 \times 105 \text{ units/mg}$.

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation:

Lyophilized from a 0.2 □m filtered concentrated solution in PBS, pH 7.4.

AA Sequence:

MAFTEHSPLTPHRRDLCSRSIWLARKL RSDLTALTESYVKHQGLNKNINLDSAD G M P V A S T D Q W S E L T E A E R L Q E N L Q A Y RTFHVLLARLLEDQQVHFTPTEGDFHQ AIHTLLLQVAAFAYQIEELMILLEYKIP RNEADGMPINVGDGGLFEKKLWGLKV LQELSQWTVRSIHDLRFISSHQTGIPAR GSHYIANNKKM

Endotoxin:

Less than 1EU/□g of rHuCNTF as determined by LAL method.

Reconstitution:

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at $\leq -20 \square C$. Further dilutions should be made in appropriate buffered solutions.

Storage:

Tel: 0086-025-86371664

Fully biologically active when compared to standard. The This lyophilized preparation is stable at 2-8 \(\text{C}\), but should be

MADE IN CHINA

Bioworld technology, co, Ltd.

No 9, weidi road Qixia District Nanjing, 210046, P, R.China.

Email: info@biogot.com Fax:0086-025-86213570

1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA. Email: info@bioworlde.com Tel: 6123263284 Fax: 6122933841

Bioworld Technology, Inc.

PRODUCT DATA SHEET

Upon reconstitution, the preparation is stable for up to one week at 2-8 \Box C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20 \Box C to -70 \Box C. Avoid repeated freeze/thaw cycles.

Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE. Made in China

1660 South Highway 100, Suite 500 St. Louis Park,MN 55416,USA. Email: info@bioworlde.com

Tel: 6123263284 Fax: 6122933841

MADE IN CHINA

Bioworld technology, co, Ltd.

No 9, weidi road Qixia District Nanjing, 210046, P, R.China. Email: info@biogot.com

Tel: 0086-025-86371664 Fax:0086-025-86213570