## PRODUCT DATA SHEET



Bioworld Technology, Inc.

# Recombinant Human Osteoprotegerin/Fc Chimera (rHuOPG-Fc)

Catalog Number: PR1106 Source:Pichia. Pastoris. Quantity:10µg/50µg/1.0mg

# **Description**

Osteoprotegerin (OPG) is a member of the TNFR superfamily that can act as a decoy receptor for RANKL. Binding of soluble OPG to sRANKL inhibits osteoclastogenesis by interrupting the signaling between stromal cells and osteoclastic progenitor cells, thereby leading to excess accumulation of bone and cartilage. OPG is expressed in a wide variety of tissues including adult heart, lung, kidney, liver, spleen, prostate, lymph node and bone marrow. OPG is secreted both as a monomeric and a dimeric protein. Its primary structure consists of seven distinct domains, four of which corresponds to the extracellular cysteine-rich domains of TNFR proteins and constitutes the soluble OPG.

## **Molecular Weight:**

Recombinant OPG/Fc contains 412 amino acid residues, including 180 residues from mature OPG (a.a 22-201) and 232 residues from the Fc protein of human IgG1, and has a calculated molecular mass of 46.5 kDa. As a result of glycosylation, the recombinant OPG/Fc migrates as a 49 kDa protein in SDS-PAGE under reducing conditions.

## **Purity:**

>90% by SDS-PAGE and HPLC analyses.

# **Biological Activity:**

Fully biologically active when compared to the standard. Determined by its ability to neutralize the stimulation of U937 cells treated with 10 ng/ml of soluble RANKL.

# **Physical Appearance:**

Sterile Filtered White lyophilized (freeze-dried) powder.

#### Formulation:

Lyophilized from a  $0.2 \square m$  filtered concentrated solution in PBS, pH 7.4.

## **AA Sequence:**

O P G 22-201: E T F P P K Y L H Y D E E T S H Q L L C DKCPPGTYLKQHCTAKWKTVCAPCPD HYYTDSWHTSDECLYCSPVCKELQYV KQECNRTHNRVCECKEGRYLEIEFCLK HRSCPPGFGVVQGTPERNTVCKRCPDG F F S N E T S S K A P C R K H T N C S V F G L L L T Q K G N A T H D N I C S G N S E S T Q K C G I D V T L F c232: EPKSSDKTHTCPPCPAPEFEGAPSV FLFPPKPKDTLMISRTPEVTCVVVDVSH EDPEVKFNWYVDGVEVHNAKTKPREE Q Y N S T Y R V V S V L T V L H Q D W L N G K E Y K CKVSNKALPTPIEKTISKAKGQPREPQV YTLPPSRDELTKNQVSLTCLVKGFYPS DIAVEWESNGQPENNYKTTPPVLDSDG S F F L Y S K L T V D K S R W Q Q G N V F S C S V M H EALHNHYTQKSLSLSPGK

#### **Endotoxin:**

Less than 1EU/□g of rHuOPG-Fc as determined by LAL method.

#### **Reconstitution:**

Tel: 0086-025-86371664

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at  $<-20\Box C$ . Further dilutions should be made in appropriate buffered so-

**MADE IN CHINA** 

Bioworld technology, co, Ltd.

No 9, weidi road Qixia District Nanjing, 210046, P, R.China.

Email: info@biogot.com Fax:0086-025-86213570

Tel: 6123263284 Fax: 6122933841

# PRODUCT DATA SHEET

lutions.

## **Storage:**

This lyophilized preparation is stable at  $2-8\Box C$ , but should be kept at  $-20\Box C$  for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at  $2-8\Box C$ . For maximal stability, apportion the recon-

stituted preparation into working aliquots and store at  $-20\Box C$  to  $-70\Box C$ . Avoid repeated freeze/thaw cycles.

# **Usage:**

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE. Made in China

Tel: 6123263284 Fax: 6122933841

**MADE IN CHINA** 

Bioworld technology, co, Ltd. No 9, weidi road Qixia District Nanjing, 210046, P, R.China.

Email: info@biogot.com
Tel: 0086-025-86371664 Fax:0086-025-86213570