

PRODUCT DATA SHEET



Bioworld Technology, Inc.

Recombinant Murine Fibroblast Growth Factor-basic (rMubFGF)

Catalog Number: PR2015

Source: Escherichia coli.

Quantity: 10µg/50µg/1.0mg

Description

FGF basic (FGF-2, HBGF-2) is one of at least 22 mitogenic proteins of the FGF family, which show 35 - 60% amino acid conservation. Unlike other FGFs, FGF acidic and basic lack signal peptides and are secreted by an alternate pathway. Storage pools within the cell or on cell surface heparan sulfate proteoglycans (HSPG) are likely. The predicted 17 kDa FGF basic isoform can be located in both the cytoplasm and the nucleus and is presumed to be the form secreted. Transcription from alternate start sites produces 21 - 24 kDa forms found only in the nucleus. High and low molecular weight human FGF basic targets the expression of different genes when expressed in NIH-3T3 cells.

Molecular Weight:

Approximately 16.2 kDa, a single non-glycosylated polypeptide chain containing 146 amino acids.

Purity:

>98% by SDS-PAGE and HPLC analyses.

Biological Activity:

Fully biologically active when compared to standard. The ED50, as determined by the dose-dependent stimulation of thymidine uptake by BaF3 cells expressing FGF receptors is <0.1ng/ml, corresponding to a specific activity of of > 1×10⁷ units/mg.

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation:

Lyophilized from a 0.2mm filtered solution in PBS, pH 7.4.

AA Sequence:

M P A L P E D G G A A F P P G H F K D P K R L Y C K N
G G F F L R I H P D G R V D G V R E K S D P H V K L Q
L Q A E E R G V V S I K G V C A N R Y L A M K E D G
R L L A S K C V T E E C F F F E R L E S N N Y N T Y R
S R K Y S S W Y V A L K R T G Q Y K L G S K T G P G
Q K A I L F L P M S A K S

Endotoxin:

Less than 1EU/mg of rMubFGF as determined by LAL method.

Reconstitution:

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at <-20°C. Further dilutions should be made in appropriate buffered solutions.

Storage:

This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. Avoid repeated freeze/thaw cycles.

Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE. Made in China

MADE IN CHINA

Bioworld Technology, Inc.

1660 South Highway 100, Suite 500 St. Louis Park, MN

55416, USA.

Tel: 6123263284

Email: info@bioworld.com

Fax: 6122933841

Bioworld technology, co, Ltd.

No 9, weidi road Qixia District Nanjing, 210046, P, R.China.

Tel: 0086-025-86371664

Email: info@biogot.com

Fax: 0086-025-86213570