

Bioworld Technology, Inc.

Recombinant Murine Interleukin-1 beta (rMulL-1β)

DescriptionDK IP VALGLKGKNLYLSCVMKDGTPTLIL-1 β is a proinflammatory cytokine produced in a variety of cells including monocytes, tissue macrophages, keratinocytes and other epithelial cells. Both IL-1 α and IL-1 β binds to the same receptor and has similar if not identical biological prop- erties. These cytokines have a broad range of activities in- cluding, stimulation of thymocyte proliferation, by inducing IL-2 release, B-cell maturation and proliferation, by inducing restaglandin and collagenase from synovial cells. However, whereas IL-1 β is a secreted cytokine, IL-1 α is predominantly a cell-associated cytokine, IL-1 α is predominantly i cells is 4 0.002 ng/ml, corresponding to a specific activity of > 5 x 108 units/mg.DK IEV A L G L K G K N L Y L S C V M K D G T P T L Q I E S V D P K K K M E K R F V P N K I E V K S K V E F E S A E F P N W Y I S T S Q A E H K P V F L G N N S G Q D II D F T M E S V S S Endotoxin: Less than 1EU/mg of mIL-1 β as determined by LAL method. Reconstitution of 0.1-10 ms/mL. Stock solutions should be the apportioned into working aliquots and store at <-20°C. For maximal stability, apportion the reconsti- tutor fluturos. Storage: This lyophilized preparation into working aliquots and store at -20°C L o-70°C. Avoid repeate

Bioworld Technology, Inc.

1660 South Highway 100, Suite 500 St. Louis Park,MN55416,USA.Email: info@bioworlde.comTel: 6123263284Fax: 6122933841

MADE IN CHINA Bioworld technology, co, Ltd. No 9, weidi road Qixia District Nanjing, 210046, P, R.China. Email: info@biogot.com Tel: 0086-025-86371664 Fax:0086-025-86213570