

PRODUCT DATA SHEET

Bioworld Technology CO., Ltd.



p-AhR (S36) Peptide

Cat No.: AP0302P

Background

The Aryl Hydrocarbon Receptor (AHR), also known as the dioxin receptor, is a ligand-activated helix/loop/helix transcription factor found in a variety of vertebrate species. The known ligands for AHR are foreign planar aromatic compounds, such as polycyclic aromatic compounds and halogenated aromatic compounds such as 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD). Unlike the steroid/thyroid hormone receptors, there is no known physiological ligand for the AH Receptor. Studies indicate that in non-ligand activated cells, AHR is found complexed with HSP90 predominantly in the cytoplasm. Upon binding to an agonist, the ligand-activated AHR is believed to transform to a nuclear, DNA binding form. This transformation process appears to involve dissociation of HSP90 from AHR followed by formation of a heterodimer with AHR nuclear translocator protein (Arnt). The AHR-ligand complex appears to initiate gene transcription of cytochrome P450 1A1.

Swiss-Prot

P35869

Applications

Blocking

Specificity

This peptide can be used with studies using AP0302 p-AhR (S36) pAb.

Purification & Purity

Synthetic peptide p-AhR (S36). (Note: the amino acid sequence is proprietary). The purity is > 98%.

Product

1 mg/ml in DI water.

Storage & Stability

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Research Use

For research use only, not for use in diagnostic procedure.