

# SLC16A5 (A193) polyclonal antibody

Catalog: AP0702 Host:

Rabbit

Reactivity: Human, Mouse, Rat

#### **BackGround:**

Monocarboxylates, such as lactate and pyruvate, play an integral role in cellular metabolism. Lactic acid is produced in large quantities as a result of glycolysis, which provides the majority of ATP to cells under normal physiological conditions. However, accumulation of lactic acid leads to a decrease in intracellular pH and cessation of glycolysis. In order for glycolysis to continue at a high rate, lactic acid must be transported out of the cell. This transport process is carried out by a family of monocarboxylate transporters (MCTs), which function as proton symports and are stereoselective for L-lactate. The MCT family consists of at least eight members, MCT1-8, which contain between 10-12 transmembrane-helical (TM) domains, with the amino and carboxy termini located in the cytoplasm. MCT6 is highly expressed in the kidneys and is thought to have a specificity for bumetanide, a loop diuretic involved in the treatment of edema. The high substrate specificity of MCT6 suggests a possible role in therapeutic drug transport and trafficking across the plasma membrane.

#### **Product:**

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

**Molecular Weight:** 

#### ~ 55 kDa

**Swiss-Prot:** 

015375

**Purification&Purity:** 

The antibody was affinity-purified from rabbit antiserum

by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

**Applications:** 

WB: 1:500~1:1000


**Storage&Stability:** 

Store at  $4 \, \mathbb{C}$  short term. Aliquot and store at  $-20 \, \mathbb{C}$  long term. Avoid freeze-thaw cycles.

### **Specificity:**

SLC16A5 (A193) polyclonal antibody detects endogenous levels of SLC16A5 protein.

#### **DATA:**



Western blot (WB) analysis of SLC16A5 (A193) polyclonal antibody at 1:500 dillution

Lane1:HEK293T whole cell lysate(40µg)

Lane2:NIH-3T3 whole cell lysate(40µg)

Lane3:PC12 whole cell lysate(40µg)

#### Note:

For research use only, not for use in diagnostic procedure.

## **Bioworld Technology, Inc.**

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA. Email: info@bioworlde.com Tel: 6123263284 6122933841 Fax:

## Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. **Email:** info@biogot.com Tel: 0086-025-68037686 0086-025-68035151 Fax: