

ULK1 (phospho-Ser636) polyclonal antibody

Catalog: AP4007

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

ULK1 and ULK2 (for UNC-51-like kinase) encode similar amino-terminal serine/threonine kinase domains, a proline/serine-rich (PS) domain, and a species conserved carboxyl-terminal domain. Both share homology with the UNC-51 kinase from *Caenorhabditis elegans* and the APG1 kinase in yeast, which are involved in axonal extension and growth, and autophagy, respectively. ULK1 and ULK2 are thought to auto-phosphorylate the PS domain in vitro, and the significant homology among vertebrates suggest that ULK1 and ULK2 are involved in the regulation of fundamental biological processes.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 150 kDa

Swiss-Prot:

O75385

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

ULK1 (phospho-Ser638) polyclonal antibody detects endogenous levels of ULK1 protein only when phosphorylated at Ser638.

DATA:



Western blot (WB) analysis of ULK1 (Phospho-Ser638) polyclonal antibody at 1:500 dilution

Lane1:L02 whole cell lysate

Lane2:L02 treated with PBS(1×PBS,PH7.4) for 1 hour whole cell lysate

Lane3:HEK293T treated with Etoposide(20ng/ml) for 6 hours whole cell lysate

Lane4:HEK293T whole cell lysate

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151