

PRODUCT DATA SHEET

Bioworld Technology CO., Ltd.



Super Vision FITC Immunofluorescence (IF) Detection System kit (Goat anti-Mouse)

Cat No.: BD5002

Introduction

Super Vision FITC Immunofluorescence (IF) Detection System is a non-biotin, one-step detection system that allows for the demonstration of antigens in paraffin-embedded tissue, cryostat sections, and cell preparations. This kit has been developed using a proprietary hyper labeling technology used to label IgG directly with more fluorescein isothiocyanate (FITC). One step polymer system provides increased sensitivity, time savings and detection simplicity. All the components contain with PBS, proteins, stabilizers and preservatives. This kit is suitable for single, double and triple immunofluorescence detect. The color is emerald green with a correct result. The FITC is easily faded with light; all experiment process need keep away from light.

Reagents

A:Blocking Reagent	10ml
B:Antibody Solution Buffer	10ml
C:Anti-Mouse IgG-FITC	50ul
D:Anti-Fading Buffer	10ml

Application

Super Vision FITC IF kit (Goat anti-Mouse) is suitable for use with mouse primary antibodies. It can apply for paraffin-embedded tissue and cryostat sections.

Storage & Shelf life

Store at 2-8 °C. Each component is stable for up to 12 Months.


Procedure

1. Deparaffinize and rehydrate tissue section;PBS/TBS wash for 2 min*3;
2. Incubate tissue in appropriate pretreatment or digestive enzyme if required for primary antibody; and PBS/TBS wash for 2 min*3;
3. Apply Blocking Reagent and incubate for 5 minutes, PBS/TBS wash for 2 min*3 (May be omitted if primary antibodies are diluted in


buffers containing normal goat serum) ;

4. Apply primary antibody and incubate according to manufacturer's recommended protocol, PBS/TBS wash for 2 min*3;
5. Add 5ul-10ul Anti-Mouse IgG-FITC to 1ml Antibody Solution Buffer and vibrate; then add that antibody solution to the sections and incubate for 30-60 min in room temperature or 37°C; PBS/TBS wash for 2 min*3 (NOTE: FITC is light sensitive. Please avoid unnecessary light exposure.);
6. Apply the Anti-Fading Buffer, and cover the microscope cover glass for observation (NOTE: It's better for observation and operation when the section is dried in no dark box before adding anti-fading buffer).

DATA



BD5002-A:IF image of GFAP (Mouse) staining in Rat brain frozen section formalin fixed tissue section. The section was then incubated with GFAP, 2.5 μ g/ml, for 12 hour 4°C and detected using BD5002. The section was mounted with Anti-fading Buffer (BD5014).



BD5002-B:IF image of NeuN (Mouse) staining in Rat brain frozen section formalin fixed tissue section. The section was then incubated with NeuN, 2.5 μ g/ml, for 12 hour 4°C and detected using BD5002. The section was mounted with Anti-fading Buffer (BD5014).

Research Use

For research use only, not for use in diagnostic procedures.

MADE IN CHINA

Bioworld Technology, Inc.

1660 South Highway 100, Suite 500 St. Louis Park,MN

55416,USA.

Email: info@bioworlde.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co, Ltd.

No 9, weidi road Qixia District Nanjing, 210046,

P, R.China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax:0086-025-68035151