

PRODUCT DATA SHEET



Bioworld Technology, Inc.

Recombinant AITRL, Mouse

Catalog Number: BK0005-50µg

Source: Escherichia coli.

Quantity: 50µg

Description:

Activation-Inducible TNF-Related Ligand (AITRL), also known as Glucocorticoid-Induced TNF-Related Ligand (GITRL), belongs to the tumor necrosis factor superfamily (TNFSF). AITRL is a Type II single transmembrane protein and shares low conservation within the extracellular domain with other TNFSF members. AITRL is expressed on macrophages, immature and mature dendritic cells and B cells. Its receptor, Activation-Inducible TNFR family Receptor (AITR), is expressed on T lymphocytes, natural killer (NK) cells, and antigen-presenting cells. After binding by AITRL, AITR can be released. AITR activation increases resistance to tumors and viral infections and is involved in autoimmune and inflammatory processes. In addition, activated AITR increases TCR-induced T cell proliferation and cytokine production and rescues T cells and NK cells from apoptosis. Recombinant mouse Activation-Inducible TNF-Related Ligand (rmAITRL) produced in E.coli is a single non-glycosylated polypeptide chain containing 127 amino acids. A fully biologically active molecule, rmAITRL has a molecular mass of 14.5 kDa analyzed by reducing SDS-PAGE and is obtained by proprietary chromatographic techniques at GenScript.

Molecular Weight:

14.5 kDa, observed by reducing SDS-PAGE.

Purity:

> 95% by SDS-PAGE and HPLC analyses.

Biological Activity:

ED50 < 5 ng/mL, measured by the amount of Interleukin-8 produced by PMBC, corresponding to a specific

activity of > 2.0 × 10⁵ units/mg.

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation:

Lyophilized after extensive dialysis against PBS.

AA Sequence:

TAIESCMVK-
FELSSSKWHMTSPKPHCVNTTSDGKLK-
ILQSGTYLIYGQVIPVDKKYIKDNAP-
FVVQIYKKNDVLQTLMNDFQILPIGGVYEL-
HAGDNIYLFKNSKDHIQKTNTYWGILMPDLPFIS

Endotoxin:

< 0.2 EU/µg, determined by LAL method.

Reconstitution:

Reconstituted in ddH₂O at 100 µg/mL.

Storage:

Lyophilized recombinant mouse Activation-Inducible TNF-Related Ligand (rmAITRL) remains stable up to 6 months at -80 °C from date of receipt. Upon reconstitution, rmAITRL remains stable up to 2 weeks at 4 °C or up to 3 months at -20 °C.

Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.