

# PRODUCT DATA SHEET



Bioworld Technology, Inc.

## Recombinant FGF-acidic, Human

Catalog Number: BK0051-10µg

Source: Escherichia coli.

Quantity: 10µg

### Description:

Fibroblast Growth Factor- acidic (FGF-acidic), also known as FGF-1 and endothelial cell growth factor, is a member of the FGF family which currently contain 23 members. FGF acidic and basic, unlike the other members of the family, lack signal peptides and are apparently secreted by mechanisms other than the classical protein secretion pathway. FGF acidic has been detected in large amounts in the brain. Other cells known to express FGF acidic include hepatocytes, vascular smooth muscle cells, CNS neurons, skeletal muscle cells, fibroblasts, keratinocytes, endothelial cells, intestinal columnar epithelium cells and pituitary basophils and acidophils. As with other FGF's, FGF acidic exhibits considerable species cross reactivity. FGF acidic and FGF basic stimulate the proliferation of all cells of mesodermal origin, and many cells of neuroectodermal, ectodermal and endodermal origin. Recombinant human Fibroblast Growth Factor-acidic (rhFGF-acidic) produced in E.coli is a single non-glycosylated polypeptide chain containing 140 amino acids. A fully biologically active molecule, rhFGF-acidic has a molecular mass of 15.8 kDa analyzed by reducing SDS-PAGE and is obtained by proprietary chromatographic techniques at GenScript.

### Molecular Weight:

15.8 kDa, observed by reducing SDS-PAGE.

### Purity:

> 95% by SDS-PAGE and HPLC analyses.

### Biological Activity:

ED50 < 0.3ng/ml, measured by a cell proliferation assay of 3T3 Cells, corresponding to a specific activity of > 3.3 × 10<sup>6</sup> IU/mg in the presence of 10 µg/ml of

heparin.

### Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

### Formulation:

Lyophilized after extensive dialysis against PBS.

### AA Sequence:

FNLPPGNYKKPKLLYCSNNGGHFL-  
RILPDGTVDGTRDRSDQHIQLQLSAESVGEVY-  
IKSTETGQYLAMDTDGLLY-  
GSQTPNEECLFLERLEEN-  
HYNTYISKKHAEKNWVGLKKN-  
SCKRGPRTHYGQKAILFLPLPVSSD

### Endotoxin:

< 0.2 EU/µg, determined by LAL method.

### Reconstitution:

Reconstituted in ddH<sub>2</sub>O or PBS at 100 µg/ml.

### Storage:

Lyophilized recombinant human Fibroblast Growth Factor- acidic (rhFGF-acidic) remains stable up to 6 months at -80 °C from date of receipt. Upon reconstitution, rhFGF-acidic should be stable up to 2 weeks at 4 °C or up to 3 months at -20 °C.

### Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.