

PRODUCT DATA SHEET



Bioworld Technology, Inc.

Recombinant ENA-78/CXCL5, Human(HEK 293-expressed)

Catalog Number: BK0302-25µg

Source: HEK 293

Quantity: 25µg

Description:

Epithelial-derived neutrophil-activating peptide 78 (ENA-78) is a small cytokine belonging to the CXC chemokine family. It is produced following stimulation of cells with the inflammatory cytokines interleukin-1 or tumor necrosis factor-alpha. Expression of ENA-78 has also been observed in eosinophils, and can be inhibited with the type II interferon, IFN- γ . ENA-78 stimulates the chemotaxis of neutrophils possessing angiogenic properties. It plays a role in reducing sensitivity to sunburn pain in some subjects, and could be a potential target used to understand more about pain in other inflammatory conditions. ENA-78 is well known to have chemotactic and activating functions on neutrophils, mainly during acute inflammatory responses. It can signal through the CXCR2 receptor. Recombinant ENA-78/CXCL5 produced in 293 cells is a single polypeptide chain containing 78 amino acids. rhENA-78/CXCL5 has a molecular mass of 8.5 kDa analyzed by reducing SDS-PAGE and is obtained by chromatographic techniques at GenScript.

Molecular Weight:

8.5 kDa, observed by reducing SDS-PAGE.

Purity:

> 98% as analyzed by SDS-PAGE.

Biological Activity:

The EC50 value of human ENA-78/CXCL5 on Ca²⁺ mobilization assay in CHO-K1/ G α 15/hCXCR2 cells (human G α 15 and human CXCR2 stably expressed in

CHO-K1 cells) is less than 200 ng/ml.

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation:

Lyophilized after extensive dialysis against PBS.

AA Sequence:

AGPAAAVLRELRCVCLQTTQGVHPK-
MISNLQVFAIGPQCSKVEVVASLK-
NGKEICLDPEAPFLKKVIQKILDGGNKEN

Endotoxin:

< 0.2 EU/µg, determined by LAL method.

Reconstitution:

Reconstituted in ddH₂O or PBS at 100 µg/ml.

Storage:

Lyophilized recombinant human ENA-78/CXCL5 remains stable up to 6 months at -80 °C from date of receipt. Upon reconstitution, human ENA-78/CXCL5 should be stable up to 1 week at 4 °C or up to 2 months at -20 °C.

Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.