

# PRODUCT DATA SHEET



Bioworld Technology, Inc.

## Recombinant TRAIL R-2, Human

Catalog Number: BK0333-50µg

Source: HEK 293

Quantity: 50µg

### Description:

TRAIL Receptor-2 is a cell-surface receptor involved in tumor necrosis factor-related apoptosis-inducing ligand (TRAIL)-induced cell-death signaling. The death ligand TRAIL bears high potential as a new anticancer agent, as binding to the death receptors TRAIL-R1 or TRAIL-R2 triggers apoptosis in most cancer cells. TRAIL-R2 is associated with a decrease in the survival rates of breast cancer patients.

### Molecular Weight:

~15 kDa, observed by reducing SDS-PAGE.

### Purity:

> 95% as analyzed by SDS-PAGE and HPLC.

### Biological Activity:

ED50 < 6 ng/ml, measured in a cell proliferation assay using RPMI-8226 cells in the presence of 25 ng/ml of human TRAIL.

### Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

### Formulation:

Lyophilized after extensive dialysis against PBS.

### AA Sequence:

ALITQQDLAPQQR AAPQQRSSPSEGLCPPGH-  
HISEDGRDCISCKYGDYS-  
THWNDLLFCLRCTRCDSEVEL-  
SPCTTTRNTVCQCEEGTFREEDSPEMCRKCR-  
TGCPRGMVKVGDCTPWSIDIECVHKE

### Endotoxin:

< 0.2 EU/µg, determined by LAL method.

### Reconstitution:

Reconstituted in ddH<sub>2</sub>O or PBS at 100 µg/ml.

### Storage:

Lyophilized recombinant Human TRAIL Receptor-2 remains stable up to 6 months at -80 °C from date of receipt. Upon reconstitution, Human TRAIL Receptor-2 should be stable up to 1 week at 4 °C or up to 2 months at -20 °C.

### Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.