

# PRODUCT DATA SHEET



Bioworld Technology, Inc.

## Recombinant TNF- $\alpha$ , Rat

Catalog Number: BK0349-50 $\mu$ g

Source: *P. pastoris*

Quantity: 50 $\mu$ g

### Description:

Tumor Necrosis Factor-Alpha (TNF-Alpha) also known as Cachectin and TNFSF1A, plays a major role in growth regulation, differentiation, inflammation, viral replication, tumorigenesis, and autoimmune diseases; and in viral, bacterial, fungal, and parasitic infections. Besides inducing hemorrhagic necrosis of tumors, TNF has been found to be involved in tumorigenesis, tumor metastasis, viral replication, septic shock, fever, inflammation, and autoimmune diseases including Crohn's disease, and rheumatoid arthritis as well as graft-versus-host disease. Recombinant rat Tumor Necrosis Factor-Alpha (rrTNF-Alpha) produced in *P. pastoris* is a glycosylated polypeptide chain containing 157 amino acids. A fully biologically active molecule, rrTNF-Alpha has a molecular mass of 17.4kDa analyzed by reducing SDS-PAGE and is obtained by proprietary chromatographic techniques at GenScript.

### Molecular Weight:

17.4kDa, observed by reducing SDS-PAGE.

### Purity:

> 95% by SDS-PAGE and HPLC analyses.

### Biological Activity:

ED50 < 50 pg/ml, measured in a cytotoxicity assay using mouse L-929 cells in the presence of actinomycin D, corresponding to a specific activity of > 2.0  $\times$  10<sup>7</sup> units/mg.

### Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

der.

### Formulation:

Lyophilized after extensive dialysis against PBS.

### AA Sequence:

MLRSSSQNSSDKPVAH-  
VVANHQAEEQLEWLSQRANALLANGMDLKDN  
QLVVPADGLYLIYSQVLFKGGQCPDYVLLTHTVS  
RFAI-  
SYQEKVSLLSAIKSPCKDTPEGAELKPWYEP-  
MYLGGVVFQLEKGDLLSAEVNLPKYL-  
DITESGQVYFGVIAL

### Endotoxin:

< 0.2 EU/ $\mu$ g, determined by LAL method.

### Reconstitution:

Reconstituted in ddH<sub>2</sub>O at 100  $\mu$ g/ml.

### Storage:

Lyophilized recombinant rat Tumor Necrosis Factor-Alpha (rrTNF-Alpha) remains stable up to 6 months at -80  $^{\circ}$ C from date of receipt. Upon reconstitution, rrTNF-Alpha should be stable up to 2 weeks at 4  $^{\circ}$ C or up to 3 months at -20  $^{\circ}$ C.

### Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.