

## PRODUCT DATA SHEET

Bioworld Technology CO., Ltd.



### Caspase 10 (K442) Peptide

Cat No.: BS1039P

#### Background

Four isoforms of caspase 10 (caspase 10a, 10b, 10c, and 10d) having the same prodomain but different mature large and small subdomain, have been described. Caspase 10 contains two death domains (DED) involved in linking to the death effector domain of the adapter protein FADD and recruiting the complex to TNFR1 and Fas. The inactive procaspase 10 is variably expressed in many tissues and cell lines as a cytosolic protein. The mature form of caspase 10 comprises two subunits, p23/p17 (splice isoforms) and p12. Interestingly, a caspase 9- dependent processing of caspase 10 by caspase 6 in cell-free extracts has recently been suggested. Caspase 10 can cleave and activate caspases 3, 4, 6, 7, 8, and 9. This is followed by cleavage of numerous key proteins, including the nuclear protein PARP.

#### Swiss-Prot

Q92851

#### Applications

Blocking

#### Specificity

This peptide can be used with studies using BS1039 Caspase 10 (K442) pAb.

#### Purification & Purity

Synthetic peptide Caspase 10 (K442). (Note: the amino acid sequence is proprietary). The purity is > 98%.

#### Product

1 mg/ml in DI water.

#### Storage & Stability

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

#### Research Use

For research use only, not for use in diagnostic procedure.