

PRODUCT DATA SHEET

Bioworld Technology CO., Ltd.



Cot (P284) Peptide

Cat No.: BS1074P

Background

The role of mitogen-activated protein kinases (MAPKs) in cell signaling pathways is well established. The rat gene Tpl-2, for tumor progression locus 2, and the human and mouse homologues c-Cot, for cancer osaka thyroid oncogene, encode a proto-oncogene serine/threonine protein kinase that was shown to play a role in the functional activation of the MAP kinase pathway. Overexpression of Cot induces MAP kinase activation in COS-1 and NIH/3T3 cells. Cot-mediated activation of MAP kinase is inhibited by both Ras N17, a dominant negative mutant of c-H-Ras, and Raf-1s621A, a dominant negative mutant of Raf-1, suggesting that Cot functions upstream of Ras and Raf-1. Other studies have shown that a kinase-negative, dominant negative mutant of Cot partially blocks Ras or Raf-1-induced MAP kinase activation, arguing that Cot functions downstream of Ras and Raf-1.

Swiss-Prot

P41279

Applications

Blocking

Specificity

This peptide can be used with studies using BS1074 Cot (P284) pAb.

Purification & Purity

Synthetic peptide Cot (P284). (Note: the amino acid sequence is proprietary). The purity is > 98%.

Product

1 mg/ml in DI water.

Storage & Stability

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Research Use

For research use only, not for use in diagnostic procedure.