

PR (V184) polyclonal antibody

Catalog: BS1299

Host: Ra

Rabbit

Reactivity: Human

BackGround:

The effects of progesterone are mediated by two functionally different isoforms of the progesterone receptor, PR-A and PR-B, which are transcribed from distinct, estrogen inducible promoters within a single copy of the PR gene. The PR-A and PR-B proteins are 90 kDa and 118 kDa respectively; the first 164 amino acids of PR-B are absent in PR-A. Progesterone bound PR-A and PR-B have different transcription activation properties. Specifically, PR-B functions as a transcriptional activator in most cell and promoter contexts, while PR-A is transcriptionally inactive and functions as a strong ligand dependent transdominant repressor of steroid hormone receptor transcriptiona activity. An inhibitory domain (ID), which maps to the amino terminus of the receptor, exists within both PR isoforms.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

- ~ 90 kDa (PR-A)
- ~ 118 kDa (PR-B)

Swiss-Prot:

P06401

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:


IHC: 1:50~1:200

Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-20 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.

Specificity:

Progesterone Receptor (V184) polyclonal antibody detects endogenous levels of Progesterone Receptor protein. DATA:



Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

 Add:
 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.

 Email:
 info@bioworlde.com

 Tel:
 6123263284

 Fax:
 6122933841

Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: <u>info@biogot.com</u> Tel: 0086-025-68037686 Fax: 0086-025-68035151