

## PRODUCT DATA SHEET

Bioworld Technology CO., Ltd.



### IKK $\beta$ (L193) Peptide

Cat No.: BS1540P

#### Background

The transcription factor NF $\kappa$ B is retained in the cytoplasm in an inactive form by the inhibitory protein I $\kappa$ B. Activation of NF $\kappa$ B requires that I $\kappa$ B be phosphorylated on specific serine residues, which results in targeted degradation of I $\kappa$ B. I $\kappa$ B kinase  $\alpha$  (IKK $\alpha$ ), previously designated CHUK, interacts with I $\kappa$ B- $\alpha$  and specifically phosphorylates I $\kappa$ B- $\alpha$  on Serines 32 and 36, the sites that trigger its degradation. IKK $\alpha$  appears to be critical for NF $\kappa$ B activation in response to proinflammatory cytokines. Phosphorylation of I $\kappa$ B by IKK $\alpha$  is stimulated by the NF $\kappa$ B inducing kinase (NIK), which itself is a central regulator for NF $\kappa$ B activation in response to TNF and IL-1. The functional IKK complex contains three subunits, IKK $\alpha$ , IKK $\beta$  and IKK $\gamma$  (also designated NEMO), and each appear to make essential contributions to I $\kappa$ B phosphorylation.

#### Swiss-Prot

O14920

#### Applications

Blocking

#### Specificity

This peptide can be used with studies using BS1540 IKK $\beta$  (L193) pAb.

#### Purification & Purity

Synthetic peptide IKK $\beta$  (L193). (Note: the amino acid sequence is proprietary). The purity is > 98%.

#### Product

1 mg/ml in DI water.

#### Storage & Stability

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

#### Research Use

For research use only, not for use in diagnostic procedure.